

wzef -!

\*Xi @'e'lb, 70D40»mgú0l/2AAyžž0i u lgp~šāÁVáC:l[ glzšá lz... ~i 0'lgzl''lwze  
 %\$cg»yÁiÆuÈgzl má lz äWDT • Nflak»l0l/2~gll f oDzZÁ[ z! á lz äW~VÇ;Š&  
 pVoi Xi μAA± ÝJc, 50~ÖÁV" 0l/2gzlk»DzZÁtÝluš~ ~šWÁ\*šJcY2050Xi Bæ°»~  
 æ'gXšèlql~ ] ášlylx • i gYK ] ášl³ , gñ" a ÆtÝl~gzlaÁwzem<lk»l0l/2Z  
 ÁVzgP»ÆwzeÆ ~k5 1Fžž0i International Program on Rice Biotechnology (IPRB)»c&fÁ  
 iÁVzgLšl0lá lz ä™i 6wzegzL+CaÆwzeÆxš6kXi ;gm'èlšælaà{pS{a Æi CQD1984  
 Xi CYÁèlšælaà • Æxš6

~T álfD ] IŠxA~ð;šÁ(80)i li ,W»] Y. \_0l' lá lz äYKĐáljÆ~g»Áwzef -!  
 wE l»V<ÖÁ) zãð0] Nqá lz äfwE l~ x explant source0cz CEB) z] gzli CEÁÁ(è 04ñ;  
 Agrobacterium gzlparticle bombardment Xá šcšmi ,Wl xj»i šZ Æ8 CŠCQ ~gzš kXá šH  
 žL>Cm0gáÉ gzl(gfp) á " zglÖ! gzl(gusA)m[čzgY%g0gÖÆáR7gÁ~ p tumefaciens-mediated  
 Xá šCwD f a Æá 3k S-IAlgzl z

wzef -!3\* ĐV-gFgzlVzQ

äàyvĀ` Áwzež@šHg»wzef -!~ Æb0gzlÄcgšÁá 3k ĐVzQ • Æxš6IPRB  
 VxÆwzeXnYÁèl0k Đ sheath blightgzl0S0bacterial blightVzgÁ>0yellow stem borer, á lz  
 f#ÿ` B, 30ã1/2DzZÁklgzli ك-ء»„ nh" Á` ~lñÆ&m<I~i0lñgzl] 0ÁVzOÆVà~  
 Šžñ0k ÁZ gšdlì 0Ā~ pool ÆáÆwzežVYi x»Ā q10k ĐVzOÆVàa ÆVzgP»Xi CY  
 a Æ0k ĐVzQ lepidopteran ~VfVšgñ" Xi ;gx»gl\*gl0q1á " hĀ0k {z`ÆkĪ 7  
 ÁwzeK! 6050kXi Ág»xi l0ÉCE6 (cry1Ab, 1Ac 1Aa, 2A, 1B, or combination of these genes)K!  
 Æ2009X^ðž™7Sg" ~gliInđĀÁwzeK! 6gī āātJc[ lèæ 0X^Á~ 1998 ~ui š«öaiWály  
 } (Ákl~ 2012 ~T ~š• ~-lĀ Bt Shanyou 63gzlBt rice Huahui No.1 ä®gi ] glizÁu ~ygzl  
 Xāi šyk»"šæe



†Yl~á 1/2

gzl(+e~Wgzl0b»0 ŽĚĪ0ēš 0ēš 0kg` Ā0ēšW) ] \*n0ð~qgzlā0" z60(\$Ā)V3g0f-gà;1g»wze  
 Ā w" »ĪgzlTV~ ~´ ~d3423Xî =gfYl»w" ĀE (CEl´ ~gzl06! ðWz0ēš0 + , ĀEg0} cā)2Wz  
 |ŠZŠ0&7p0V"gz\$ÐzĀkXî CyfŋĀVb07%ēl~V-š!WylÐzĀT0i @YH{Sci¹ wE l»wzei  
 b) °»~b7ĀVfzšj ] gzlĪG02°YXî CyfµzŋĀZĪgzšÑ0~0} lðWz~VĀúnqgz&pàzäö  
 ĀVfV½x• i ĩ ĩf:æWš 2™fē. t6x b76ylgzlĪfµ»] cšZy0i šžñx° Ā' Je~šā~Tī @7t™o, »  
 Tī i šqĪ»YB(-It0i i ĩ ĩ x b76»عكس g~l~T0i ;gYHwE l»g»i šh}a ĀE±Yl~iG02  
 ðWz6wzef -İĐáĪj kXî @Yf†Yl™fē. ‡~iG02Ā` ÐzĀX• DY0Š™4šā, l~V" ~  
 àzŋŋŠgzla{Sci} gfĀEöŰlkwĀ` ÐšāĀ™gā~c&lgzl÷š@0i Ywzet25 24i šHj»(ōgyōĪ)} l  
 °@ĐáĪjĀē™~gYJY2012x ĪZ ĀwzeyōĪXī šHj») gfĀEwpĀEloci" ZPg~±òĀĀwze  
 ~@gzl Ī%ā~yĪšcī ĀVfa/ŠkiZĀwzef -li qĀzbēš9°YXī ~gYx»6• z68i'ei™gā  
 ~½\*š) gfĀE±YĀEYlzbēš9~šÑĀEōžī Qe³Ā] tklyĪÐk0cW~} @xtYl{C" ~gĪ ĀšÑ  
 Xī YYHI ~i gZ0»ZĀEŋĀšÑ

ĀE±YĀEzbēš9~wzeXDYñ07 í tēš gZ0 ðñ0@ēR0i t-ē} CEĪĀā0iñ%o~wze  
 ~R šziWĀēš9NDØ& ~OZ ĪE- ðñ) gfĀERNAi approachžlfX¥tÐ] !šā lz äYKĐáĪj  
 ĪĪzi VÈ] It 27XH†Yl~i, gĪE(CElSg3lgzĪ t-ē} CEĪĀĪY Ð3ĪgāĀE- ðñā klgzllf†Yl  
 ) gfĀE†Yl~gĀE" z6k0i Cf: xa ĀEVāšVzgaYgzlVāšl" z6àz äfYqÐVzš7~«ĀE  
 Xī µĀĀfi XēlqĪ½\*š

Biopharming ÐWze

àz ä3àz äYĀg»ÐwzeXī i ĩ ĩ ē z~Tī @YHwE l ~~g»Ā] cšZĀXĪ LZ Āwze  
 yX• DYf! ~] Yx4zhĀE^ Ā5qĀEYžž0i cholera toxin B subunit (CTB) XĀlqĪ~ē z  
 H4šB, ĀEòx~@Ā; šg0 ĪmucosalM° a ĀEXĀž@ @YHŸqÐ] YmucosalĀE~Wgzl} nĀ  
 Ā klgzl Cf 7 [ b76] gĪwzšĀE} #JĀ²ZĪ Ðw, |ĪeCTB {š™g»Ðwze {z~ĀEkl 28XnY  
 gzl} nē z-wšĀ mucosal cellXī Sg0ÐxĀWēš9~wjā; i tgzlXī SgĪk@immunogenicity

q{ZĀa ĀŠĀlā z• g ~ ˘ 7k5 1FĐZĀäf: t0• C- s@ĀV-gFàz äf ~ 5lgĀ ÷ , gzläv  
Xī Cf{Sci} gzcā V;žVY• ŠfU

laØk ~ žL>ŠgpZá z äYñ0 ~ VâĀlgzVwñ0VzġŌV2%wE {SciĐu} cšS-!A  
~ yJĀEwzeiLiġŌŋz+9} 2qġiS-!A%Xi ~ gYI Āi wš •ĀES-!AĐZĀT Ōi eš™  
%ñî wzei qĀEĀ ū iLiġŌŋz+9} ġi C!ŠĀV2%b'Xi „ gYĀXaġĀä™a) gfĀEYB(-!  
c]legzlj ĀVŌĀi Ō kIXlfĲŸL~ òx bacteriostaticgzltg• LCġE; ĀVŌŌ &Ō%ĀyġDT  
Xa ĀEV"ŠBñĀEVâĀlgzVzġĀYLgm<li YYHW E la ĀefXĐ

wgzlwzef -!

iGġĀwzeŌtŸL~ «o) gfĀEŠŠĐf!ŠS-!} lgzŌfXĐV-gFgzVzQ~ Š»Āwzef -!  
wzef -!Xī YYHW E !Wġ• ~ ylyĀ~i, ] cšĀĀk{z ĀEKĲi ġtŸL~Ĳ ĲĀ ) gfĀEĲŸL~  
~Šā%ĐáJĀE ĀwzeB, B, ĀEKIXi YO»1 zġĀtŸLK-ĀE] gġwzš) gfĀE~4~ ] 1j ā  
Yl»ä™sp~ [, WŌle Oġ»gzlŌtĲĀĀŸzġĀĲz{ " CžĲi ġtŸL wš» C4 Pathway ā ~ X• ~gY] ġ  
ò}VĲlylŌ•ĀEKlgzĲi [fi, W»i ~ŠāĲ ĐáJĀEgzlaĀVœy³cwzeapomict{z ĀEKĲŌi =gf  
Šæ~ ä™y»i Ō 5ĐáJĀE%ĀwzegzĲifμzŲŲcū~ ±ŲĲ~gzlaĀwzeo'ĐkĲIX• %ofqzŲfĀ  
XīA

:] Y!j

Rice Biotechnology: A need for developing countries. 2004 Swapan K Data Agbioforum 7(1&2):31-35. 141210  
Rice Biotechnology: Rockefeller to End Network After 15 Years of Success 1999. Dennis Normille 2  
(<http://www.sciencemag.org/>)  
IRRI Annual Report, 1999. Los Baños, Laguna 3  
Achieving successful deployment of Bt Rice. 2004. Sasha Ming High, Michael B. Cohen, Qing Yao Shu, and 4  
Illimar Altosaar. TRENDS in Plant Science 9:6:286-292.  
Translational fusion hybrid Bt genes confer resistance against yellow stem borer in transgenic elite 5  
Vietnamese rice (Oryza sativa L.) cultivars. 2006. N. H. Ho, N. Baisakh, N. Oliva, K, Datta, R. Frutos, and S.K. Datta.  
Crop Scie. 46:781-789

Bt rice harbouring cry genes controlled by a constitutive or wound-inducible promoter: protection and transgene expression under Mediterranean field conditions. 2004. J.C. Breitler, J.M. Vassal, Maria del Mar Catala, D. Meynard, V. Marfa, E. Mele, Monique Royer, Isabel Murillo, Blanca San Segundo, E. Guiderdoni, and Joaquina Messeguer. *Plant Biotechnology Journal* 2(5) :417-430. 6

Transgenic rice plants with a synthetic cryIAbi gene from *Bacillus thuringiensis* were highly resistant to eight lepidopteran rice pest species. Q. Shu, G. Ye, H. Cui, X. Chang, Y. Xiang, D. Wu, M. Gao, Y. Xia, C. Hu, R. Sardana, and I. Altosaar. *Mol. Breed.* 6: 433-439. 7

Transgenic IR72 with fused Bt gene cry1(b)/cry1A(c) from *Bacillus thuringiensis* is resistant against four lepidopteran species under field conditions. 2001. *Plant Biotechnol.* 18:125-133. 8

Crop Biotech Update 4 December, 2009. 9

A receptor kinase-like protein encoded by the rice disease resistance gene Xa21. 1995. Song, W.Y. Song, G.L. Wang, L.L. Chen, H.S. Kim, L.Y. Pi, T. Holsten, B. Wang, W.X. Zhai, H. Zhu, C. Fauquet, and P.C. Ronald, *Science* 270:1804-1806. 10

Transgenic elite indica rice varieties, resistant to *Xanthomonas oryzae* pv. *Oryzae*. 1998. S Zhang, W-Y. Song., L. Chen, D-L. Ruan, N. Taylor, P. Ronald, R. Beachy and C. Fauquet *C. Molecular Breeding* 4:551-558. 11

Field performance of Xa21 transgenic indica rice (*Oryza sativa* L.), IR72. 2000. *Theoretical and Applied Genetics.* 101:15-20. 13

Genetic engineering of rice for resistance to sheath blight. 1996. W. Lin, C. Anuratha, S. Datta, K. Potrykus, S. Mathukrishnan and S.K. Datta. 1995. *Bio/Technology* 13:686-691. 15

Improving rice to meet food and nutrient needs: Biotechnological approaches. 2002 S.K. Datta and G.S. Khush. *Journal of Crop Production.* 6(1):229-247. 16

The map-based sequence of the rice genome. 2005. International Rice Genome Sequencing Project. *Nature,* Vol. 436, pp. 793-800. 17

<http://www.agbios.com/dbase.php?action=ShowProd&data=CL121%2C+C1141%2C+CFX51> 18

Water management in irrigated rice: Coping with water scarcity. 2007. B.A.M. Bouman, R.M. Lampayan, and T.P. Tuong. [http://dspace.irri.org:8080/dspace/bitstream/10269/266/2/9789712202193\\_content.pdf](http://dspace.irri.org:8080/dspace/bitstream/10269/266/2/9789712202193_content.pdf) 19

Improvement of water use efficiency in rice by expression of HARDY, an Arabidopsis drought and salt tolerance gene. 2007. A. Karaba, S. Dixit, R. Greco, A. Aharoni, K.R. Trijatmiko, N. Marsch-Martinez, A. Krishnan, K. N. Nataraja, M. Udayakumar, and A. Pereira. *PNAS.* 104(39):15270-15275. 20

Arabidopsis CBF3/DREB1A and ABF3 in transgenic rice increased tolerance to abiotic stress without stunting growth. 2005. S-J. Oh, S.I. Song, Y.S. Kim, H-Y. Jang, S.Y. Kim, M. Kim, Y-K. Kim, B.H. Nahm, J-K Kim. *Plant Physiology* 138:341-351. 21

Expression of a bifunctional fusion of the Escherichia coli genes for trehalose-6-phosphate synthase and trehalose-6-phosphate phosphatase in transgenic plants increases trehalose accumulation and abiotic stress tolerance without stunting growth. 2003. I-C. Jang, S-J. Oh, J-S. Seo, W-B. Choi, S.I. song, S.H. Kim, Y.S. Kim, H-S Seo, Y.D.

Choi, B.H. Nahm, and J-K. Kim. Plant Physiology. 131:516-524.

Food composition and nutrition tables. 6th ed. 2000. S.W. Souci, W. Fachmann, H. Kraut. Food Quality Control, World Food Programme. <http://foodquality.wfp.org/FoodProcessing/Cerealmilling/Nutritionalimpact/tabid336/PageContentID/458/Default.aspx>.

Engineering the provitamin A (b-carotene) biosynthetic pathway into carotenoid-free) rice endosperm. Y. Xudong, S. Al-Babili, A. Klöti, J. Zhang, P. Lucca, P.Beyer, I. Potrykus. Science 287:303-305.2000,

Improving the nutritional value of Golden Rice through increased pro-vitamin A content. 2004. J.A. Paine, C. A. Shipton, S. C. Rhian, M. Howells, M. J. Kennedy, G.Vernon, S.Y. Wright, E. Hinchliffe, J. L. Adams, A. L. Silverstone, and R. Drake. Nature Biotechnology 23(4), 482-487

Transgenic rice is a source of iron for iron-depleted rats. 2002. L.E. Murray-Kolb, F. Takaiwa, F. Goto, T. Yoshihara, E. C. Theil, and J.L. Beard. Journal of Nutrition.132:957-960.

Regulation of lysine synthesis and catabolism in rice. 2008. Q-Q. Liu, M-L. Chan, R-X. Duan, H-X. Yu, M-H. Gu, S.S. M. Sun. Abstract in Plant Genomics in China, PGCIX. (<http://www.plantgenomics.cn/abslist.cgi?absid=789>).

Rice-based mucosal vaccine as a global strategy for cold-chain and needle-free vaccination. 2007. T. Nochi, H. Takagi, Y. Yuki, L. Yang, T. Masumura, M. Mejima, U. Nakanishi, A. Matsumura, A. Uozumi, T. Hiroi, S. Morita, K. Tanaka, F. Takaiwa, H. Kiyono. PNAS. 104(26):10986-10991.

When expressing lactoferrin and lysozyme has antibiotic-like properties when fed to chicks. 2002. B.D. Humphrey, N. Huang, K.C. Klasing. J. Nutrition. 132:1214-1218.

The rice squad. 2002. Nature 416: 576-578. April 11, 2002.

Apomixis: The plant breeder's dream. 2001. (<http://www.grain.org/seedling/?id=2>)

International Service for the Acquisition of Agri-biotech Applications (ISAAA) SEAsiaCenter c/o IRRI, DAPO Box 7777,

X, TM, jDÍ Y

Tel: +63 2 845 0563

Fax: +63 2 845 0606

E-mail:[knowledge.center@isaaa.org](mailto:knowledge.center@isaaa.org)





ìl4l 6 q1XŠ cz™ Sg" ~ xn Agrobacterium tumefaciens strain CP4  
žŽĀĀá lz äfwE la Ægzla Ā ^ h} Ā 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS)  
gzl%Ā} Š7ì gŠ) f»; @Ā] · %gŠÒp} uzŠgzli (E1ĀgŠÒp3hð6 CŠq10i @™x» ~ 5 hshikimate  
• „ ÜxI lî lzg] : SĀxn~ig\l&fjžì \_f"U] ItĐ ] Y. \_óžpšXŠc0iĀa Æ ] Š  
ØXì H7ê »ä™76äeCg^ ~ L gâJlĀ ĀKl ä Monsantoë@

w »xnf -!

### xn3& ĐVzhc} Q

(Tilletia ōí x-àz• ~ V[ gzl ŽSA0(Puccinia graminis), P~ Tì @fo, »ōí ĀxI lFĀxn  
žŽì ōí [ lyĐf ŽSA ~ ð; ylx • DY' Đ~! ~ ` Āxn~ iŠŽñĀ&t • i ōí tritici)  
la 7 äŠÌ ÇIQc• Dfla äŠ Lgð• l~ # • ~ ĒÆKl0 • سن » head blight & crown rot  
0° 0...Ā• z » 1/2 ĀVzgāYcVāc{lZ>gzl• DfYCiQc• D- Ğ+IĀðāžD>%oÆ ŽSAXDf  
1āHqzĀx»6" μÆä™g»xn3& Đ ŽSAä SyngentaX • سن » V-gF-uzŠgzl! ly~ yiĀÆyñg;  
äfla ĐáljÆxnf -IXŠcŠ™yñĐzĀ] Ýl< lá lz äYK6YB (-ID±YĀxlĀ" μKl~ Y2007  
~g»Āxn3& Đōí àz• ~ Vā0 Syngenta ōXì Šfglylo Çq1x ā9& Fusarium Ā7 ÆpŠDqàz  
lXÇñYHwE l»ú yzç™gāa ÆÑ K0i „ g™x»pB, ÆCIMMYTA Æ

Š~xn) gfÆ ] Y. \_{e{eä > Æ6'W4lyYgzl^ kXlYBĒ Ğöi†~ cg-i ~ Y2009  
Āā Lr34 ä > Æcg-i03Xì Š™7i + 4»} çÆōí ~ xni t0Hs'ì l»iŠŽñĀā 3& Đōí  
ì @Yc0 ~ n1 ĀxnžŽHĀcğš ā Yr36 ä VāŠb , Æç 4lyY • D™kç S ÜÆV-gFZ žŽĀh ,  
gzlĀ @™ç ÜĀā Šñ~ ~gFYr36žŃi³ tĐi Klä VāŠb , Xì @Yf#Ÿ~ çÆö0l6Rwyp  
Xì @™-gYe@ĀŠg3&

xn ĀxI lZ {zXì Ā~g»tāu~ (Đf~ ylyÆxnf -lä(CAAS)b , x~vls Wò07i  
Yellow X • i ō9 3& ĐVzhc} Qgzl-ōí Ā ^ ĀS` 0head scab0K]zyellow mosaic ~ T • i ™g»  
nĀxnàz ä™ĀĀ;ÆUYyĀ~ ; gE- x~vly | 05l4Xíf [ 0š ~ gzi lJY2015n3& Đ Mosaic Virus  
76gî Cg^ JY2013cY2012ntĀxnXnYHĀĀyv , 20á lz äfĐzĀäWyĀ~ Đ0%ž@ „ gYĀg»  
l6Xì ñY~Š™

xnàzä™"Š'Ä#

t(Nax1 and Nax2) • a™Äcgs, - ÇZÄEná z äYÄ# ä ^ a^g~7&l-öCISRO  
Vp+gzlaa Ä} \$7žž òì HÖl»ŽÎä ÇVäSX • t ÐTriticum monococcum Ý ÐnäbÄxn  
ÄíÑá Nax2 ~ ] !Šályá z äYK ~ ´7W Y2009 Ç7X • MfÇUYC i™wle ^ zyg ~ 5gyxgÄ  
X~Šgzla {Sci, 25 ~ iŠŽñÄe gðä Vžš

ì ÇfÐzzÄWE IÄgluten~gFtòifpôa ÄŠbZgD»V-gFÄùžž • „gYÄg»x l+lÄxn  
d zXi @YfgD»ŦÄi Gb! %ÐzzÄT òì rle ^ zyg ~ x ÇVÄYl bð½q(ZÄE™ [ lyÄ-WKggz  
gliadins á z ä™la-gFÄùÐ ~ xn~Tì „gYÄg»xn+l ñfD™WE IÄeÇCQ~ ; gE-Ä l  
qtì µÄäf {Sci} gll Ä- ðÑ~nkXifµz~4h~ ~ α SÄÖÄxnÐ klgzÇñYcšwiC!Ä  
Ç8Xi @Yc0Ä¹ ~ xnžžì ÇEÄäòìÑ

xnàzä™"Š'ä, È

X +FèlqŦÄä0ÐzzÄ~pŠñ@~gžš{Šžñm<li !pèlqŦÄáJÄxnØ& Ðà, È  
ò, ÄÄV; zÐzzÄà, È ~ ´7W} òì C™Ä%] ZWZ} • 6u Äoì ÈŦ~gzlaÄ` Çl Ç9Xi „gYK  
Ä, 70~ðäÄxnÐzzÄà, È h" ~Y2007 ~tgÍzX(ylgzÄEY2003ÐY2002) Ŧ~fµzŦÄ, qŦ~äæW  
Wz6Äm7&l-Îl6SW¹ g6e~tgÍz~Y2007XZ7Ç™"Š' {gÄ}°le • 300Ä, ÇgÐzzÄT òŦ~fµzŦ  
thale cress (Zea mays), h~T ~ íÑCÄxnXK ] !ŠóíÑ è ÇR 30Äxnä Spangenberg Ö  
Äà, È • i Ç (Saccharomyces cerevisiae). P(Physcomitrella-patens) ð»(Arabidopsis thaliana)  
Ä pathways ð6 ÇŠžž • D™ encode ~È×gÄ" z6+lâtxñfÄcgs ä 6 á z ä™Š'Š'  
ià, È ...~VB, &{ÖWÄ~f [ x»rtÄxnÇXi BìiÄ%òÄÄ} Š7l ~gll ÄÄä0 ÄE™ regulate  
ä International Maize and Wheat Improvement Center (CIMMYT) bŠìlÇXì ñYf ½xnàz äY  
8b lCžXì ÄÖÄÄÄ™†Ýl~ ] : SÄxnñfD™WE l(DREB1A) äqŦÄ Arabidopsis thaliana  
Ç1Xi i qÄÇ&Ää™"Š'Äe gðz] gmwzŠÄ0à, È xnt{Š™g»Ð

µñlžšÄ Äxn f -!

ä™76gî Çg~Äyl~Y2004gzY2003ÐzzÄ~p~ÝÄ~gàa Äxn f -lç(GM) {" ÇfCQ  
ä™"»Äxnä(NAWG) ^ ò, àz äÇlxn Ç2Xi Ç #ðf f(ÄV" è ÇRzÄk lŠcŠ™yñÄ" µÄ



Fusarium Head Blight and Mycotoxin Contamination of Wheat, A Review.1990. CHA Snijders. European Journal of Plant Pathology. DOI: 10.1007/BF01974256. 10

USDA FORUM: Syngenta Glad to See Support for Biotech Wheat. 2010. Dow Jones Newswires. <http://www.nasdaq.com/aspx/company-news-story.aspx?storyid=201002181922dowjonesdjonline000783#ixzz0v2tw09s3> 11

Syngenta Foundation for Sustainable Agriculture launches Syngenta and CIMMYT Wheat Ug99 Stem Rust Resistance Research Partnership. 2009. Syngenta Foundation for Sustainable Agriculture. <http://www.syngentafoundation.org/db/1/821.pdf>. 12

Biotech Could Save World Wheat Crops. 2009. Committee for A Constructive Tomorrow. <http://www.cfact.org/a/1527/Biotech-could-save-world-wheat-crops>. 13

China approves big budget for GMO amid food worries. 2008. N. Shuping. <http://www.reuters.com/article/environmentNews/idUSPEK11727520080710?feedType=RSS&feedName=environmentNews&sp=true> 14

China Plans \$3.5 Billion GM Crops Initiative. 2008. R. Stone. Science. DOI: 10.1126/science.321.5894.1279. 15

Expression of Endogenous Thioredoxin Gene and  $\alpha$ -Amylase Activity in Transgenic Wheat with Antisense-Thioredoxin Gene at Grain Filling Stage. 2007. R. Jiang-ping, et al. Journal of Triticeae Crops. DOI: 27(5):893-897. 16

Physiological Characterization of Two Genes for Na<sup>+</sup> Exclusion in Durum Wheat, Nax1 and Nax2. 2006. RA James, et al. Plant Physiology. DOI: 142:1537-1547. 17

GM Wheat Means Hope for Celiac Sufferers. 2010. A. Nemeth. <http://www.foodsafetynews.com/2010/01/genetically-modified-foods-are-becoming-1/>. 18

Drought-tolerant Wheat: Promising Results. 2008. GMO Safety. <http://www.gmo-safety.eu/science/grain/583.drought-tolerant-wheat-promising.html> 19

Results of Transgenic Wheat Trial Look Promising. 2004. CIMMYT Newsletter. <http://www.cimmyt.org/index.php/en/newsletter/120-2004/391-results-of-transgenic-wheat-trial-look-promising>. 20

22 Monsanto Pulls the Plug on Genetically Modified Wheat. 2004. Science Magazine. DOI: 10.1126/science.304.5674.1088a. 21

76% of Wheat Growers Approve Biotech Petition. 2009. National Association of Wheat Growers. <http://www.wheatworld.org/wp-content/uploads/biotech-petition-press-release-20090226.pdf>. 22

2010 Consumer Perceptions of Food Technology Survey. 2010. International Food Information Council. [http://www.foodinsight.org/Content/3843/Final\\_Executive%20Summary%20Food%20Tech%20Report\\_Website%20version\\_7-7-10.pdf](http://www.foodinsight.org/Content/3843/Final_Executive%20Summary%20Food%20Tech%20Report_Website%20version_7-7-10.pdf) 23

Will Cleverly Crafted Prose Win Over Science and Evidence? The Case of GM Wheat. 2010. P Fitzgerald. Agrifood Awareness Australia Limited. [http://www.afa.com.au/letters\\_editor/GM\\_wheat\\_prose\\_or\\_science.pdf](http://www.afa.com.au/letters_editor/GM_wheat_prose_or_science.pdf). 24

The Case for Biotech Wheat. 2009. Wheat Foods Council. 26

<http://www.wheatfoods.org/FileLibrary/FileImage/FINAL%20The%20Case%20for%20Biotech%20Wheat.pdf>

X • DàJ \ W ~ i 0 U - qzly, VĀ] à ¥ O D Y B ( - I Ž Ž • a Ā d { S M # • D á j A E Y B ( - I Ā V " K = 0

International Service for the a A E ] à ¥ n \* X ( <http://www.isaaa.org/kc> ) X i H j » ä 0 % @ - A E Y B ( - I Ā V " Ā V " Ā y l

0 3 0 z 0 Acquisition of Agri-biotech Applications (ISAAA) SEAsiaCenter c/o IRRI, DAPO Box 7777,

X, ™, 0 D í Y

Tel: +63 2 845 0563

Fax: +63 2 845 0606

E-mail: [knowledge.center@isaaa.org](mailto:knowledge.center@isaaa.org)

# xnf -!

VÆKX • i Š ū àl™i ~: WäŠÆTì V { " çfqtì ð7ÑzpgmDy10{Æk vxn  
 »«ztægjî ð5l' Xì @Y'gcl'9MÄ{ á lz ä 3Æ` tlgz • DfñfñÈÐ } uδq1~ ] gBÁÚpq1  
 ~ ~g»Ä) zDŠgzl0600~. 00KzgMwtgzlì @YÑJe ~ ^ ÄLWÄTì ` tLq1xn Xì ölañYÄxn' ~  
 ~ xI làz äYðçLÄxnÄ` WXì @YHwE l~ÄÄ3AÄ0gzl2 ] lzæÆ [ NÄkLz`ÆKXì @fWE l  
 äfWE l~ äz) Triticum durumgzl(xnàz äfWE l~ Kzg) Triticum aestivum L.XI lX- {Šci Df  
 X • (xnàz

la xn~ \*Š5X ¶Hu• 683.15gzla ~ Y2009ÄKXì ` tLÑz äfWE l{Šci Df xn^ Äh  
 Äo xnàz äflaV(Epì oL(DfÑz ä™la xnu X • i pM%0c&0u ~ ~ } {&á lz ä™  
 (1 i e™la xn:Ñ, ý<sup>hE</sup> t 2D1.3w, CM%Xì e™la xn. \_Æ ] gzçKlì c&Xì CYfWE l, g0L  
 Klì í o glgz1 7W0" Xì @Yc™æW o yZÜÜ »Klp bushel of wheat at 13.5% moisture=27.21kg)  
 6Xì ;gf Ä w" »S l@ÆKlgzì ðfµz ¶~gzla ÄxnylgzŠÆð;Š, \*X • i Š™æW o yZÜz q1»  
 zÄg»i Šî lzg á lz äYKg (la Äö0) LÄxnžVYì gD»: gzla Äxn~ « ÄÄÊ cîgzlh  
 gzi6ä™g (lj Sh) ÄEYB (-la Äw" Ägzla °YÄxn [ pVOçXì ú' ÐV" Váz {gÄetÐ  
 Xì e tÐj Šî lzg t™i ÄX • MYKi b) {zÐŠæÄTì ;gYcŠ

## xnî qÄ' ð' žD

ì clearfield wheat Xç»T 0ðz™sg" 6äeCg ^ xnàz ð' žD » ~Y2007 ~l" äBASf  
 lz žD tì Sg{0i1 ~ iŠžñÄ imidazolinone herbicide žžì gzla Äö0) l{ " çf Clearfield wheatX  
 ~ 5 lg dç» SÄCElÄÄlgŠçp ~gzç Ca Ä%Ä} Š7AHASXì ÄÄöuÄ (AHAS) çç (EÄ Wzge; I+l  
 gzla Ä00%Äé ~ YžÄò) {Š™Y qÐöäi wälz äYÄÐ á jÄKl~M%Xì i lz^ Nž äW  
 6Xì /xÄè) L lclearfield Ðá jÄyDÄö0) LÄk1~ agzLØk ÐV-gF0(á)  
 ~ Äg» • ÄMonsanto, the MON 71800) gfÄ8 ççÄE CO` «Äxnàz ä™ ð' žD Cm  
 ŽÜ x-çq1Nž äYc0 ~ } ia Ää™g» xnàz ð' žD Ä ðtXì Roundup Ready WheatXç Cg ^ »Kl

ìl4l 6 q1XŠ cz™ Sg" ~ xn Agrobacterium tumefaciens strain CP4  
žŽĀĀā á lz äfwE la ĀgZla Ā ^ h} Ā 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS)  
gzl%Ā} Š7i gŠ) f»; @Ā] · %gŠ Āp} uzŠgzli (E1ĀgŠ Āp3 hđ6 CŠq10i @™x» ~ 5 hshikimate  
• „ ÜxI lî lzg] : SĀxn~i g\l&f hžī \_f"U] ItĐ ] Y. \_óžpđXŠc0iĀa Ā ] Š  
0Xī H7ē »ä™7 b äeCg^ ~ L gâJlĪ ĪĀkl ä Monsantoë@

w »xnf -!

### xn3& ĐVzhc} Q

(Tilletia 0ī x-āz• ~ V[ gzl ŽSA0(Puccinia graminis), P~ Tī 0fo, »0ī ĀxI lFĀxn  
žŽī 0ī [ lyĐf ŽSA ~ đ; ylx • DY' Đ~! ~ ` Āxn~i ŠŽñĀ&t • i 0ī tritici)  
la 7 äŠ Ī ÇIQc• Dfla äŠ Lgđ• l~ # • ~ ĒĀKl0 • سن » head blight & crown rot  
0° 0...Ā• z » 1/2 ĀVzgāYcVāc{lZ>gzl• DfYCiQc• D- Ą ĪĀđāžD>%oĀ ŽSAXDf  
1āHqzĀx»6" μĀä™g»xn3& Đ ŽSAä SyngentaX • سن » V-gF-uzŠgzl! ly~ yiĪĀEyñg;  
äfla Đá lĀxnf -IXŠcŠ™yñĐzĀ] Ÿl< lá lz äYK6YB (-ID± YĀxlĪ" μkl~ Y2007  
~g»Āxn3& Đ0ī āz• ~ Vā0 Syngenta 0Xī Šfglylo Cq1x āđ9& Fusarium Ā7 ĀpŠDqāz  
0XÇñYHwE l»ú yzç™gāa ĀĪ K0ī „ g™x»pB, ĀCIMMYTA Ā

Š~xn) gfĀ ] Y. \_{e{e ä > Ā6đ'W4lyYgzl^ kXlYBĀĀ öi†~ cg-i ~ Y2009  
Āā Lr34 ä > Ācg-i03Xī Š™7i + 4»} çĀ0ī ~ xni t0Hs Ī l»i ŠŽñĀā 3& Đ0ī  
i @Yc0 ~ n1 ĀxnžŽHĀcğš ā Yr36 ä VāŠb , Āç 4lyY • D™k S ÜĀEV-gFZ žŽĀh ,  
gzlī @™ġ ĪĪŠñ~ ~gFYr36žŃi³ tĐi klä VāŠb , Xī @Yf#Ÿ~ ĄĀö0l Ī6Rwyp  
Xī @™-gYe@ĀŠg3&

xn ĀxI lZ {zXī Ā~g»tāu~ (Đf~ ylyĀxnf -lä(CAAS) b , x~vls Wb0lī  
Yellow X • i 9 3& ĐVzhc} Qgzl-0ī Ā ^ ĀS` 0head scab0k]zyellow mosaic ~ T • i ™g»  
nĀxnāz ä™ĀĪĄĀUYyĀ~ ; gE- x~vly | 0504Xīf [ 0š ~ gzi ĪĪY2015n3& Đ Mosaic Virus  
7 6gī Cg^ JĪY2013cY2012ntĀxnXnYHĀĀyv , 20á lz äfĐzĀäWyĀ~ Đ0%ž@ „ gYĀg»  
06Xī ĩY~Š™

xnàzä™"Š'Ä#

t(Nax1 and Nax2) • a™Äcgs, - ÇZÄXná lz äYÄ# ä ^ a^g~7&L-öCISRO  
Vp+gzlaa Ä} \$7žž òì HÖl»ŽÎä ÇVäSX • t ÐTriticum monococcum Ý ÐnäbÄxn  
ÄíÑá Nax2 ~ ] !Šályá lz äYK ~ ´7W Y2009 Ç7X • MfÇUYC i™wle ^ zyg ~ 5gyxgÄ  
X~Šgzla {Sci, 25 ~ iŠžÑÄe g0ä Vžš

ì ÇfÐzzÄWE IÄgluten~gFt0íf pöa ÄŠb lgD»V-gFÄüžž • „gYÄg»x l+lÄxn  
d lzXì @YfgD»¶Ä i Gb! %ÐzzÄT òì rle ^ zyg ~ x ÇÄÄYl b0½q(ZÄE™ [ lyÄ-WKggz  
gliadins á lz ä™la-gFÄüÐ ~ xn~Tì „gYÄg»xn+l ñfD™WE IÄeGÇQ~ ; gE-Ä l  
qtì µÄÄä f {Sci} gll Ä - ðÑ ~ nkXíf µlz~4h<sup>2</sup> ~ α SÄÖÄxnÐ klgzÇñYcšwiC!Ä  
Ç8Xì @Yc0Ä¹ ~ xnžžì ÇEÄäiòìÑ

xnàzä™"Š'ä, È

X +Fèlq¶¶Ää0ÐzzÄ~pŠñ@~gžš{Šžñm<li !pèlq¶Ðá jÄxn0& Ðà, È  
ò ÄÄV;žÐzzÄà, È ~ ´7W} òì C™Ä%] ZWZ} • 6u Äoì È¶~gzlaÄ` Çl Ç9Xì „gYK  
Ä, 70~ðäÄxnÐzzÄà, È h" ~ Y2007 ~tgÍzX(ylgžšÄY2003ÐY2002) ¶fµz¶Ä, q¶~äæW  
Wz6Äm7&l-Îl6SW¹ g6e~tgÍz~Y2007XZ7Ç™"Š' {gÄ}°le • 300Ä, ÇgÐzzÄT 0¶fµz¶¶  
thale cress (Zea mays), h~T ~ íÑCÄxnXK ] !ŠóíÑ è ÇR 30Äxnä Spangenberg Ö  
Äà, È • i ¶ (Saccharomyces cerevisiae). P0(Physcomitrella-patens) ð»0 (Arabidopsis thaliana)  
Ä pathways ð6 Çšžž • D™ encode ~È<sup>2</sup>gÄ" z6+lâtxñfÄcgs ä 6 á lz ä™Š'Š'  
iXì ä, È ...~VB, &{ÖWÄ~f [ x»rtÄxnÇXì Bì iÄ%òÄÄ} Š7l ~gll ÄÄä0 ÄE™ regulate  
ä International Maize and Wheat Improvement Center (CIMMYT) bŠì lÇXì ñYf ½xnàz äY  
gb lCžXì ÄÖÄÄä™¶Ýl~ ] : SÄxnñfD™WE l(DREB1A) äq¶Ä Arabidopsis thaliana  
Ç1Xì i qÄÇ&Ää™"Š'Äe g0z] gmwžšÄ0à, È xnt{Š™g»Ð

µñlžšÄ Äxn f -!

ä™76gî Çg^Äyl~Y2004gzY2003ÐzzÄ~p~ÝÄ~gàa Äxn f -lç(GM) {" ÇfÇQ  
ä™"Š'»Äxnä(NAWG) ^ ò, àz äÇlxn Ç2Xì Ç #ðf f (ÄV" è ÇRzÄk lŠcŠ™yñÄ" µÄ



Fusarium Head Blight and Mycotoxin Contamination of Wheat, A Review.1990. CHA Snijders. European Journal of Plant Pathology. DOI: 10.1007/BF01974256. 10

USDA FORUM: Syngenta Glad to See Support for Biotech Wheat. 2010. Dow Jones Newswires. <http://www.nasdaq.com/aspx/company-news-story.aspx?storyid=201002181922dowjonesdjonline000783#ixzz0v2tw09s3> 11

Syngenta Foundation for Sustainable Agriculture launches Syngenta and CIMMYT Wheat Ug99 Stem Rust Resistance Research Partnership. 2009. Syngenta Foundation for Sustainable Agriculture. <http://www.syngentafoundation.org/db/1/821.pdf>. 12

Biotech Could Save World Wheat Crops. 2009. Committee for A Constructive Tomorrow. <http://www.cfact.org/a/1527/Biotech-could-save-world-wheat-crops>. 13

China approves big budget for GMO amid food worries. 2008. N. Shuping. <http://www.reuters.com/article/environmentNews/idUSPEK11727520080710?feedType=RSS&feedName=environmentNews&sp=true> 14

China Plans \$3.5 Billion GM Crops Initiative. 2008. R. Stone. Science. DOI: 10.1126/science.321.5894.1279. 15

Expression of Endogenous Thioredoxin Gene and  $\alpha$ -Amylase Activity in Transgenic Wheat with Antisense-Thioredoxin Gene at Grain Filling Stage. 2007. R. Jiang-ping, et al. Journal of Triticeae Crops. DOI: 27(5):893-897. 16

Physiological Characterization of Two Genes for Na<sup>+</sup> Exclusion in Durum Wheat, Nax1 and Nax2. 2006. RA James, et al. Plant Physiology. DOI: 142:1537-1547. 17

GM Wheat Means Hope for Celiac Sufferers. 2010. A. Nemeth. <http://www.foodsafetynews.com/2010/01/genetically-modified-foods-are-becoming-1/>. 18

Drought-tolerant Wheat: Promising Results. 2008. GMO Safety. <http://www.gmo-safety.eu/science/grain/583.drought-tolerant-wheat-promising.html> 19

Results of Transgenic Wheat Trial Look Promising. 2004. CIMMYT Newsletter. <http://www.cimmyt.org/index.php/en/newsletter/120-2004/391-results-of-transgenic-wheat-trial-look-promising>. 20

22 Monsanto Pulls the Plug on Genetically Modified Wheat. 2004. Science Magazine. DOI: 10.1126/science.304.5674.1088a. 21

76% of Wheat Growers Approve Biotech Petition. 2009. National Association of Wheat Growers. <http://www.wheatworld.org/wp-content/uploads/biotech-petition-press-release-20090226.pdf>. 22

2010 Consumer Perceptions of Food Technology Survey. 2010. International Food Information Council. [http://www.foodinsight.org/Content/3843/Final\\_Executive%20Summary%20Food%20Tech%20Report\\_Website%20version\\_7-7-10.pdf](http://www.foodinsight.org/Content/3843/Final_Executive%20Summary%20Food%20Tech%20Report_Website%20version_7-7-10.pdf) 23

Will Cleverly Crafted Prose Win Over Science and Evidence? The Case of GM Wheat. 2010. P Fitzgerald. Agrifood Awareness Australia Limited. [http://www.afa.com.au/letters\\_editor/GM\\_wheat\\_prose\\_or\\_science.pdf](http://www.afa.com.au/letters_editor/GM_wheat_prose_or_science.pdf). 24

The Case for Biotech Wheat. 2009. Wheat Foods Council. 26

<http://www.wheatfoods.org/FileLibrary/FileImage/FINAL%20The%20Case%20for%20Biotech%20Wheat.pdf>

X • DàJ \ W ~ i 0 U - qzly, VĀ] à ¥ O D Y B ( - I Ž Ž • a Ā d { S M # ĩ Đ á j A E Y B ( - I Ā V " K = 0

International Service for the a A E ] à ¥ ĩ \* X ( <http://www.isaaa.org/kc> ) X ĩ H j » ä 0 % @ - A E Y B ( - I Ā V " Ā V " Ā y l

0 3 0 z 0 Acquisition of Agri-biotech Applications (ISAAA) SEAsiaCenter c/o IRRI, DAPO Box 7777,

X, ™, ĩ Đ í Ÿ

Tel: +63 2 845 0563

Fax: +63 2 845 0606

E-mail: [knowledge.center@isaaa.org](mailto:knowledge.center@isaaa.org)

wzef -!

\*Xi @'e'lb, 70D40»mgú07/2ÁÁyžž0i u lgp~šāÁVáC:l[ glzšá lz... ~i 0'lgzl''lwze  
 %\$cg»yÁiÆuÈgzl má lz äWDT • Nflak»l07/2~gll f oDzZÁ[ z! á lz äW~VÇ;Š&  
 pVoi Xi μÁ± ÝJc, 50~ÖÁV" 07/2gzlk»DzÁtÝluš~ ~šWÁ\*šJcY2050Xi Bæ°»~  
 æ<sup>E</sup>gXšèlq[~ ] ášlylx • i gYK ] ášl<sup>3</sup>, gñ" a ÆtÝl~gzlaÁwzem<lk»l07/2Z  
 ÁVzgP»ÆwzeÆ ~k5 1Fžž0i International Program on Rice Biotechnology (IPRB)»c&fÁ  
 iÁVzgšl0lá lz ä™i 6wzegz+CaÆwzeÆx<sup>6</sup>6KXi ;g™èššæläá{pS{a Æi CQD1984  
 Xi CYÁèššæläá• Æx<sup>6</sup>6

~T álfD ] IŠxA~ð;šÁ(80)i li ,W»] Y. \_0' lá lz äYKĐáljÆ~g»Áwzef -!  
 wE l»V<ÖÁ) zãð0] Nqá lz äfwE l~ x explant source0cz CB) z] gzli CEÁá(è 04ñ;  
 Agrobacterium gzlparticle bombardment X á šcšmi ,Wì xj»i šZ Æ8<sup>E</sup>lcQ ~gzšKXá šH  
 žD>Cm0<sup>E</sup>gáÉ gzl(gfp) á " z6glÖ! gzl(gusA)m[čzgY%g0gÖÆáR7gÁ~ p tumefaciens-mediated  
 Xá šCBñD f a Æá 3k S-IAlgzl z

wzef -!3\* ĐV-gFgzVzQ

äàyvÃ` Áwzež@šHg»wzef -!~ Æb0'gzlÄcgšÁá 3k ĐVzQ• Æx<sup>6</sup>6IPRB  
 VxÆwzeXnYÁè0k Đ sheath blightgzl0S0bacterial blightVzgÁ>0yellow stem borer, á lz  
 f#ÿ` B, 30ã1/2DzÁklgzli ك-ء»„ nh" Á` ~lñÆ&m<I~i0ñgzl] 0ÁVzOÆVà~  
 Šžñ0k ÁZgšdlì 0Á~ pool ÆáÆwzežVYi x»Á q10k ĐVzOÆVàa ÆVzgP»Xi CY  
 a Æ0k ĐVzQ lepidopteran ~VfVšgñ" Xi ;gx»gl\*gl0q1á " hÁ0k {z`ÆKlì 7  
 ÁwzeK! 6050KXl Ág»xI l0ÉCE6 (cry1Ab, 1Ac 1Aa, 2A, 1B, or combination of these genes)K!  
 Æ2009X^ðž™7Sg" ~gliIn0ÁÁwzeK! 6gî äätJc[ lèæ 0X^Á~1998 ~ui š«öaiWály  
 } (Ákl~2012~T~š• ~-lÁ Bt Shanyou 63gzlBt rice Huahui No.1 ä@gi ] glizÁu ~ygzl  
 Xái pyk»"»æäe



†Yl~á 1½

gzl(+e~Wgzl0b»0 ŽĚĪ0ēš 0ēš 0kg` Ā0ēšW) ] \*n0ð~gZlā# " z60(\$Ā)V3g0f-gà;1g»wze  
 Ā w" »~ÍgzlTV~ ~´ ~d³423Xì =gfYl»w" ĀE (CEl´ ~gzl06! ðWz0ēšt0 + , ĀEg0} cā)2Wz  
 |ŠZŠ0&7p0V"gz\$ÐzZĀkXì CyfŋĀVb0½ēl~V-Š!WylÐzZĀT 0i @YH{Šci¹ wE l»wzei  
 b) °»~b7ĀVfzšj ] gZlĪG½°YXì CyfµzŋĀZĪgzšÑ0~0} lðWz~VĀúnqzL&pàz äö  
 ĀVfV½• i ĩ ĩ tf: æWš 2™fē. t6× t6yLgzlĪfµ»] cšZy0iŠžñ° Ā' Je~šā~Tì @7t™o, »  
 Tì i ŠqĪ»YB(-It0i ĩ ĩ ĩ xBz6»عكس g~l~T 0i ;gYHwE l»g»i Šh}a ĀE±Yl~ĪG½  
 ðWz6wzef -lÐáĪj kXì @YftYl™fē. ‡~ĪG½Ā` ÐzZĀX• DY0Š™4šā, l~V" ~  
 àzŋĪŠgzla{Šci) gfĀEöŰlkwĀ` ÐšāĀ™gā~c&lgz÷\$@0i Ywzet25 24i ŠHj»(ōgyōÍ)} l  
 °@ĪÐáĪjĀē™~gYJY2012X ĪZ ĀwzeyōÍXì ŠHj») gfĀEwpĀEloci" ZPg~±òĀĀwze  
 ~@gzl Ī%ā~yĪĀšcī ĀVfa/ŠkiZĀwzef -li qĀEzbēš9°YXì ~gYx»6• z68i'ei™gā  
 ~½\*š) gfĀE±YĀEYlzbēš9~šÑĀEōžī Qe³Ā] tKlylÐkZ0cW~} @xtYl{C" ~gĪ ĀšÑ  
 Xì YYHI ~i gZ0»ZĀEŋĀšÑ

ĀE±YĀEzbēš9~wzeXDYñ07 ĩ tēgZl0 ðñ0@ēR0i t-ē} CEĪĀl0iñ%o~wze  
 ~R šziWĀēš9NDØ&~0Z ĪE- ðñ) gfĀERNAi approachžlfX¥tÐ] !Šá lz äYKÐáĪj  
 ~Ízi VÈ] It 27XHtYl~i, gĪE(CElSg3lgzĪ t-ē} CEĪĀlY Ð3ĪgāĀE- ðñā klgzllftYl  
 ) gfĀE†Yl~gĀE" z6kZ0i Cf: xa ĀEVāšVzgaYgzlVāšl" z6àlz äfYqÐVzš7~«ĀE  
 Xì µĀĀfi XēlqĪ½\*š

Biopharming ÐWze

àz ä3àz äYĀg»ÐwzeXì ĩ ĩ ĩ ē z~Tì @YHwE l ~g»Ā] cšZĀXĪ LZ Āwze  
 yX• DYf! ~] Yx4zhĀE^ Ā5qĀEYžž0i cholera toxin B subunit (CTB) XĀlqĪ~ē z  
 H4šB, ĀEòu~@Ā; Šg0 ĪmucosalM° a ĀEXĀž@ @YHŸqÐ] YmucosalĀE~Wgzl} nĀ  
 Ā klgzL Cf 7 [ 7y6] gĪwzšĀE} #JĀá²ZĪ Ðw, |ĪeCTB {Š™g»Ðwze {z~ĀEkl 28XnY  
 gZl} nē z~wšĀ mucosal cellXì ŠgpÐxĀWēš9~wjā; ĩ tgzXì ŠĪk@immunogenicity

q{ZĀa ĀŠĀlā z• g ~ ˆ 7k5 1FĐZĀäf: t0• C- s@ĀV-gFàz äf ~ 5lgĀ ÷ , gzläv  
Xī Cf{Sci] gzcā V;žVY• ŠfU

laØk ~ žL>ŠgpZá z äYñ0 ~ VâĀlgzlvwñ0VzġĠ0V2%wE L{SciĐu} cšLŠ-!A  
~ yJĀEwzeiLiĠđŃgzl+9<sup>đđ</sup> } 2đđ S-!A%Xi ~ gYl Āi wš •ĀES-!AĐZĀT Ōi eš™  
%ñġi wzei qĀEĀ Ōi LiĠđŃgzl+9<sup>đđ</sup> ġi C!ŠĀV2%b'Xi „ gYĀXaĠĀä™<sup>a</sup>) gfĀEYB(-!  
c]legzlj ĀVŌĀi Ō KIXlfĲYl~ đđ bacteriostaticgzltg• LCĀEj ĀVŌŌ &đđ%ĀYĠĐT  
Xa ĀEV"ŠBñĀEVâĀlgzlvzġĀYLgm<li YYHwE la ĀefXĐ

wgzlvzef -!

iGġĀwzeŌtYl~ «o) gfĀEŠ-Đf!ŠS-!} lgzġfXĐV-gFgzlvzQ~ Š»Āwzef -!  
wzef -!Xī YYHwE !Wġ• ~ ylyĀ~i, ] cšĀĀklz ĀEKĲi ġtYl~Ġ LĀ ) gfĀEĲYl~  
~Šā%ĐáJĀE ĀwzeB, B, ĀEKIXi YO»1 zġĀtYlK-ĀE] gġwzš) gfĀE~4~ ] 1j ā  
Yl»ä™sp~ [, Wđle OġgzlđtĲĀĀYzġĀZ{" ČZĲi ġtYl wš» C4 Pathway ā ~ X• ~gY] ġ  
ò}YĲlyđ•ĀEKlgzli [fi, W»i ~ŠāĲ ĐáJĀEgzlaĀVœy<sup>3</sup>cwzeapomict{z ĀEKĲŌi =gf  
Šæ~ ä™y»i Ō 5ĐáJĀE%Āwzegzli fμzŲVcú~ đđ~gzlzaĀwzeo' ĐKĲIX• %ofqđđfĀ<sup>a</sup>  
XīA

:] Y!j

Rice Biotechnology: A need for developing countries. 2004 Swapan K Data Agbioforum 7(1&2):31-35. 141210  
Rice Biotechnology: Rockefeller to End Network After 15 Years of Success 1999. Dennis Normille 2  
(<http://www.sciencemag.org/>)  
IRRI Annual Report, 1999. Los Baños, Laguna 3  
Achieving successful deployment of Bt Rice. 2004. Sasha Ming High, Michael B. Cohen, Qing Yao Shu, and 4  
Illimar Altosaar. TRENDS in Plant Science 9:6:286-292.  
Translational fusion hybrid Bt genes confer resistance against yellow stem borer in transgenic elite 5  
Vietnamese rice (Oryza sativa L.) cultivars. 2006. N. H. Ho, N. Baisakh, N. Oliva, K, Datta, R. Frutos, and S.K. Datta.  
Crop Scie. 46:781-789

Bt rice harbouring cry genes controlled by a constitutive or wound-inducible promoter: protection and transgene expression under Mediterranean field conditions. 2004. J.C. Breitler, J.M. Vassal, Maria del Mar Catala, D. Meynard, V. Marfa, E. Mele, Monique Royer, Isabel Murillo, Blanca San Segundo, E. Guiderdoni, and Joaquina Messeguer. *Plant Biotechnology Journal* 2(5) :417-430. 6

Transgenic rice plants with a synthetic cryIAbi gene from *Bacillus thuringiensis* were highly resistant to eight lepidopteran rice pest species. Q. Shu, G. Ye, H. Cui, X. Chang, Y. Xiang, D. Wu, M. Gao, Y. Xia, C. Hu, R. Sardana, and I. Altosaar. *Mol. Breed.* 6: 433-439. 7

Transgenic IR72 with fused Bt gene cry1(b)/cry1A(c) from *Bacillus thuringiensis* is resistant against four lepidopteran species under field conditions. 2001. *Plant Biotechnol.* 18:125-133. 8

Crop Biotech Update 4 December, 2009. 9

A receptor kinase-like protein encoded by the rice disease resistance gene Xa21. 1995. Song, W.Y. Song, G.L. Wang, L.L. Chen, H.S. Kim, L.Y. Pi, T. Holsten, B. Wang, W.X. Zhai, H. Zhu, C. Fauquet, and P.C. Ronald, *Science* 270:1804-1806. 10

Transgenic elite indica rice varieties, resistant to *Xanthomonas oryzae* pv. *Oryzae*. 1998. S Zhang, W-Y. Song., L. Chen, D-L. Ruan, N. Taylor, P. Ronald, R. Beachy and C. Fauquet *C. Molecular Breeding* 4:551-558. 11

Field performance of Xa21 transgenic indica rice (*Oryza sativa* L.), IR72. 2000. *Theoretical and Applied Genetics.* 101:15-20. 13

Genetic engineering of rice for resistance to sheath blight. 1996. W. Lin, C. Anuratha, S. Datta, K. Potrykus, S. Mathukrishnan and S.K. Datta. 1995. *Bio/Technology* 13:686-691. 15

Improving rice to meet food and nutrient needs: Biotechnological approaches. 2002 S.K. Datta and G.S. Khush. *Journal of Crop Production.* 6(1):229-247. 16

The map-based sequence of the rice genome. 2005. International Rice Genome Sequencing Project. *Nature,* Vol. 436, pp. 793-800. 17

<http://www.agbios.com/dbase.php?action=ShowProd&data=CL121%2C+C1141%2C+CFX51> 18

Water management in irrigated rice: Coping with water scarcity. 2007. B.A.M. Bouman, R.M. Lampayan, and T.P. Tuong. [http://dspace.irri.org:8080/dspace/bitstream/10269/266/2/9789712202193\\_content.pdf](http://dspace.irri.org:8080/dspace/bitstream/10269/266/2/9789712202193_content.pdf) 19

Improvement of water use efficiency in rice by expression of HARDY, an Arabidopsis drought and salt tolerance gene. 2007. A. Karaba, S. Dixit, R. Greco, A. Aharoni, K.R. Trijatmiko, N. Marsch-Martinez, A. Krishnan, K. N. Nataraja, M. Udayakumar, and A. Pereira. *PNAS.* 104(39):15270-15275. 20

Arabidopsis CBF3/DREB1A and ABF3 in transgenic rice increased tolerance to abiotic stress without stunting growth. 2005. S-J. Oh, S.I. Song, Y.S. Kim, H-Y. Jang, S.Y. Kim, M. Kim, Y-K. Kim, B.H. Nahm, J-K Kim. *Plant Physiology* 138:341-351. 21

Expression of a bifunctional fusion of the Escherichia coli genes for trehalose-6-phosphate synthase and trehalose-6-phosphate phosphatase in transgenic plants increases trehalose accumulation and abiotic stress tolerance without stunting growth. 2003. I-C. Jang, S-J. Oh, J-S. Seo, W-B. Choi, S.I. song, S.H. Kim, Y.S. Kim, H-S Seo, Y.D.

Choi, B.H. Nahm, and J-K. Kim. Plant Physiology. 131:516-524.

Food composition and nutrition tables. 6th ed. 2000. S.W. Souci, W. Fachmann, H. Kraut. Food Quality Control, World Food Programme. <http://foodquality.wfp.org/FoodProcessing/Cerealmilling/Nutritionalimpact/tabid336/PageContentID/458/Default.aspx>.

Engineering the provitamin A (b-carotene) biosynthetic pathway into carotenoid-free) rice endosperm. Y. Xudong, S. Al-Babili, A. Klöti, J. Zhang, P. Lucca, P.Beyer, I. Potrykus. Science 287:303-305.2000,

Improving the nutritional value of Golden Rice through increased pro-vitamin A content. 2004. J.A. Paine, C. A. Shipton, S. C. Rhian, M. Howells, M. J. Kennedy, G.Vernon, S.Y. Wright, E. Hinchliffe, J. L. Adams, A. L. Silverstone, and R. Drake. Nature Biotechnology 23(4), 482-487

Transgenic rice is a source of iron for iron-depleted rats. 2002. L.E. Murray-Kolb, F. Takaiwa, F. Goto, T. Yoshihara, E. C. Theil, and J.L. Beard. Journal of Nutrition.132:957-960.

Regulation of lysine synthesis and catabolism in rice. 2008. Q-Q. Liu, M-L. Chan, R-X. Duan, H-X. Yu, M-H. Gu, S.S. M. Sun. Abstract in Plant Genomics in China, PGCIX. (<http://www.plantgenomics.cn/abslist.cgi?absid=789>).

Rice-based mucosal vaccine as a global strategy for cold-chain and needle-free vaccination. 2007. T. Nochi, H. Takagi, Y. Yuki, L. Yang, T. Masumura, M. Mejima, U. Nakanishi, A. Matsumura, A. Uozumi, T. Hiroi, S. Morita, K. Tanaka, F. Takaiwa, H. Kiyono. PNAS. 104(26):10986-10991.

When expressing lactoferrin and lysozyme has antibiotic-like properties when fed to chicks. 2002. B.D. Humphrey, N. Huang, K.C. Klasing. J. Nutrition. 132:1214-1218.

The rice squad. 2002. Nature 416: 576-578. April 11, 2002.

Apomixis: The plant breeder's dream. 2001. (<http://www.grain.org/seedling/?id=2>)

International Service for the Acquisition of Agri-biotech Applications (ISAAA) SEAsiaCenter c/o IRRI, DAPO Box 7777,

X, TM, jDÍ Y

Tel: +63 2 845 0563

Fax: +63 2 845 0606

E-mail:[knowledge.center@isaaa.org](mailto:knowledge.center@isaaa.org)

